

## Riferimenti bibliografici

1. Kuwahara T, Asanami S, Kubo S. *Experimental infusion phlebitis: tolerance osmolality of peripheral venous endothelial cells*. Nutrition 1998;14(6): 496–501.
2. Isaacs JW, Millikan WJ, Stackhouse J, Hersh T, Rudman D. *Parenteral nutrition of adults with 900-milliosmolar solution via peripheral vein*. American Journal of Clinical Nutrition 1977;30:552–9.
3. Hoffmann E. *A randomised study of central venous versus peripheral intravenous nutrition in the perioperative period*. Clinical Nutrition 1989;8(4):179–80.
4. Kane KF, Cologiovanni L, McKiernan J, Panos MZ, Ayres RC, Langman MJ, et al. *High osmolality feedings do not increase the incidence of thrombophlebitis during peripheral I.V. nutrition*. Journal of Parenteral and Enteral Nutrition 1996;20(3):194–7.
5. Matsusue S, Nishimura S, Koizumi S, Nakamura T, Takeda H. *Preventive effect of simultaneously infused lipid emulsion against thrombophlebitis during postoperative parenteral nutrition*. Surgery Today 1995;25(8):667–71.
6. Timmer JG, Schipper HG. *Peripheral venous nutrition: the equal relevance of volume load and osmolarity in relation to phlebitis*. Clinical Nutrition 1991;10(2):71–5.
7. Palmer D, McFie J. *Venous access for parenteral nutrition*. In: Payne James J, Grimble GK, editors. *Artificial nutrition support in clinical practice*. London: Silk DBA; 2001. p. 379–400.
8. Ryder M. *Evidence-based practice in the management of vascular access devices for home parenteral nutrition therapy*. Journal of Parenteral and Enteral Nutrition 2006;30:S82–93.
9. Raad I, Davis S, Becker M, Hohn D, Houston D, Umphrey J, Bodey GP, et al. *Low infection rate and long durability of nontunneled silastic catheters. A safe costeffective alternative for long-term venous access*. Archives of Internal Medicine 1993;153:1791–6.
10. Ryder M. *Peripheral access options*. Surgical Oncology Clinics of North America 1995;4:395–427.
11. Safdar N, Maki DG. *Risk of catheter-related bloodstream infection with peripherally inserted central venous catheters used in hospitalized patients*. Chest 2005;128:489–95.
12. Macdonald S, Watt AJB, McNally D, Edwards RD, Moss JG, et al. *Comparison of technical success and outcome of tunneled catheters inserted via the jugular and subclavian approaches*. Journal of Vascular and Interventional Radiology 2000;11:225–31.
13. Pittiruti M, Buononato M, Malerba M, Carriero C, Tazza L, Gui D, et al. *Which is the easiest and safest technique for central venous access? A retrospective survey of more than 5,400 cases*. Journal of Vascular Access 2000; 1: 100–7.
14. Pittiruti M, Migliorini I, Emoli A, et al. *Preventing central venous catheter related infections: catheter site selection and insertion technique significantly affect the chances of adequate catheter site care*. 20th Annual Congress European

Society of Intensive Care Medicine, Berlin, 2007, Intensive Care Medicine Suppl. Sept 2007, p. S13.

15. Randolph AG, Cook DJ, Gonzales CA, Pribble CG. *Ultrasound guidance for placement of central venous catheters: a meta-analysis of the literature*. Critical Care Medicine 1996; 24 (12): 2053–8.

16. Rothschild JM, The AHRQ Committee. *Making health care safer a critical analysis of patient safety practices. Evidence report/technology assessment: number 43*. AHRQ, [www.ahrq.gov/clinic/ptsafety/summary.htm](http://www.ahrq.gov/clinic/ptsafety/summary.htm); July 2001.

17. Keenan SP. Use of ultrasound to place central lines. Journal of Critical Care 2002;17(2):126–37.

18. Hind D, Calvert N, McWilliams R, Davidson A, Paisley S, Beverley C, Thomas S, et al. *Ultrasonic locating devices for central venous cannulation: meta-analysis*. BMJ 2003;327 (7411): 361.

19. Calvert N, Hind D, McWilliams R, Davidson A, Beverley CA, Thomas SM, et al. *Ultrasound for central venous cannulation: economic evaluation of cost-effectiveness*. Anaesthesia 2004;59 (11): 1116–20.

20. Leung J, Duffy M, Finckh A. *Real-time ultrasonographically-guided internal jugular vein catheterization in the emergency department increases success rates and reduces complications: a randomized, prospective study*. Annals of Emergency Medicine 2006;48(5):540–7.

21. Karakitsos D, Labropoulos N, De Groot E, Patrianakos AP, Kouraklis G, Poularas J, et al. *Real-time ultrasound-guided catheterisation of the internal jugular vein: a prospective comparison with the landmark technique in critical care patients*. Critical Care, <http://ccforum.com/content/10/6/R162>, 2006;10:R162.

22. LeDonne J. *The age of reason: the end of blind sticking*. Annual Congress of the Association for Vascular Access, Phoenix, 2007. Handout on [www.avainfo.org](http://www.avainfo.org).

23. Parkinson R, Gandhi M, Harper J, Archibald C. *Establishing an ultrasound guided peripherally inserted central catheter (PICC) insertion service*. Clinical Radiology 1998;53:33–6.

24. Simcock L. *No going back: advantages of ultrasound guided upper arm PICC placement*. JVA 2008;13 (4): 191–7.

25. Pittiruti M, Scoppettuolo G, Emoli A, et al. *Parenteral nutrition through ultrasound- placed PICCs and midline catheters is associated with a low rate of complications: an observational study*. Nutritional Therapy and Metabolism, in press.

26. Povoski SP. *A prospective analysis of the cephalic vein cutdown approach for chronic indwelling central venous access in 100 consecutive cancer patients*. Annals of Surgical Oncology 2000; 7: 496–502.

27. Silberzweig JE, Sacks D, Azita S, The members of the Society of Interventional Radiology Technology Assessment Committee. *Reporting standards for central venous access*. Journal of Vascular and Interventional Radiology 2003; 14: S443–52.

28. Francis KR, Picard DL, Fajardo MA, Pizzi WF. *Avoiding complications and decreasing costs of central venous catheter placement utilizing*

*electrocardiographic guidance. Surgery Gynecology and Obstetrics* 1992;175:208–11.

29. Pittiruti M, Scoppettuolo G, LaGreca A, et al. *The EKG method for positioning the tip of PICCs: results from two preliminary studies.* *JAVA* 2008; 13(4): 179–86.

30. Maki DG, Kluger DM, Crnich CJ. *The risk of bloodstream infection in adults with different intravascular devices: a systematic review of 200 published prospective studies.* *Mayo Clinic Proceedings* 2006;81(9): 1159–71.

31. Hockenhull JC, Dwan K, Boland A, et al. *The clinical and cost effectiveness of central venous catheters treated with anti-microbial agents in preventing bloodstream infections: a systematic review and economic evaluation.* *Health Technology Assessment*, www.hta.nhsweb.nhs.uk; 2006.

32. Dezfulian C, Lavelle J, Nallamothu BK, Kaufman SR, Saint S, et al. *Rates of infection for single-lumen versus multilumen central venous catheters: a metaanalysis.* *Critical Care Medicine* 2003;31:2385–90.

33. Zürcher M, Trämer M, Walder B. *Colonization and bloodstream infection with single- versus multi-lumen central venous catheters: a quantitative systematic review.* *Anesthesia and Analgesia* 2004;99:177–82.

34. Garnacho-Montero J, Aldabero S, Palla S T, Palomar-Martínez M, Valles J, Almirante B, Garces R, et al. *Risk factors and prognosis of catheter related bloodstream infection in critically ill patients: a multicenter study.* *Intensive Care Medicine* 2008;34:2185–93.

35. Turcotte S, Dube S, Beauchamp G. *Peripherally inserted central venous catheters are not superior to central venous catheters in the acute care of surgical patients on the ward.* *World Journal of Surgery* 2006; 30: 1605–19.

36. Deshpande KS, Hatem C, Ulrich HL, Currie BP, Aldrich TK, Bryan-Brown CW, Kvetan V, et al. *The incidence of infectious complications of central venous catheters at the subclavian, internal jugular, and femoral sites in an intensive care unit population.* *Critical Care Medicine* 2005; 33(1).

37. Lorente L, Villegas J, Martin MM, Jimenez A, Mora ML, et al. *Catheter-related infection in critically ill patients.* *Intensive Care Medicine* 2004; 30: 1681–4.

38. Durbec O, Viviani X, Potie F, Vialet R, Albanese J, Martin C, et al. *A prospective evaluation of the use of femoral venous catheters in critically ill adults.* *Critical Care Medicine* 1997; 25: 1986–9.

39. Reusch S, Walder B, Tramer MR. *Complications of central venous catheters: internal jugular versus subclavian access – a systematic review.* *Critical Care Medicine* 2002; 30: 454–60.

40. Hamilton HC, Foxcroft DR. *Central venous access sites for the prevention of venous thrombosis, stenosis and infection in patients requiring long-term intravenous therapy.* *Cochrane Database of Systematic Reviews* 2007; (3). doi:10.1002/14651858.CD004084.pub2. Art. No.: CD004084.

41. Warren DK, Zack JE, Mayfield JL, Chen A, Prentice D, Fraser VJ, et al. *The effect of an education programme on the incidence of central venous catheter associated bloodstream infection in a medical ICU.* *Chest* 2004; 126: 1612–8.

42. East D, Jacoby K. *The effect of a nursing staff education program on compliance with central line care policy in the cardiac intensive care unit.* Pediatric Nursing 2005; 31: 182–4.
43. Pronovost P, Needham D, Berenholtz S, Sinopoli D, Chu H, Cosgrove S, et al. *An intervention to decrease catheter-related bloodstream infections in the ICU.* The New England Journal of Medicine 2006; 355: 2725–32.
44. Boyce JM, Pittett D. HICPAC committee and HICPAC/SHEA/APIC/IDSA *Hand Hygiene Task Force: guideline for hand hygiene in health-care settings.* MMWR Recommendations and Reports 2002;51(RR16):1–44.
45. Mimoz O, Villeminey S, Ragot S, Dahyot-Fizel C, Laksiri L, Petitpas F, Debaene B, et al. *Chlorhexidine-based antiseptic solution vs alcohol-based povidone-iodine for central venous catheter care.* Archives of Internal Medicine 2007; 167 (19): 2066–72.
46. Hanazaki K, Shingu K, Adachi W, Miyazaki T, Amano J, et al. *Chlorhexidine dressing for the reduction in microbial colonization of the skin with central venous catheters: a prospective randomized controlled trial.* Journal of Hospital Infection 1999; 42: 165–7.
47. Maki DG, Mermel LA, Kluger D, et al. *The efficacy of a chlorhexidine-impregnated sponge for the prevention of intravascular-related infections: a prospective, randomized, controlled, multicenter study.* Washington, DC: American Society for Microbiology; 2000.
48. Ho KM, Litton E. *Use of chlorhexidine-impregnated dressing to prevent vascular and epidural catheter colonization and infection: a meta-analysis.* Journal of Antimicrobial Chemotherapy 2006; 58: 281–7.
49. Casey AL, Worthington T, Lambert PA, Quinn D, Farouqi MH, Elliott TS, et al. *A randomized, prospective clinical trial to assess the potential infection risk associated with the PosiFlow needleless connector.* Journal of Hospital Infection 2003; 54: 288–93.
50. McKee R, Dunsmuir R, Whitby M, Garden OJ. *Does antibiotic prophylaxis at the time of catheter insertion reduce the incidence of catheter-related sepsis in intravenous nutrition?* Journal of Hospital Infection 1985; 6: 419–25.
51. Ranson MR, Oppenheim BA, Jackson A, Kamthan AG, Scarffe JH. *Double-blind placebo controlled study of vancomycin prophylaxis for central venous catheter insertion in cancer patients.* Journal of Hospital Infection 1990; 15: 95–102.
52. Ljungman P, Hägglund H, Björkstrand B, Lönnqvist B, Ringdén O. *Peroperative teicoplanin for prevention of gram-positive infections in neutropenic patients with indwelling central venous catheters: a randomized, controlled study.* Support Care Cancer 1997;5:485–8.
53. Siegman-Igra Y, Anglim AM, Shapiro DE, Adal KA, Strain BA, Farr BM, et al. *Diagnosis of vascular catheter-related bloodstream infection: a meta-analysis.* Journal of Clinical Microbiology 1997; 35: 928–36.
54. Blot F, Nitenberg G, Chachaty E, Raynard B, Germann N, Antoun S, et al. *Diagnosis of catheter-related bacteremia: a prospective comparison of the time to positivity of hub blood versus peripheral-blood cultures.* Lancet 1999; 354: 1071–7.

55. Raad I, Hanna HA, Alakech B, Chatzinikolaou I, Johnson MM, Tarrand J. *Differential time to positivity: a useful method for diagnosing catheter-related bloodstream infections*. Annals of Internal Medicine 2004; 140 (1): 18–25.
56. Li JS, Sexton DJ, Mick N, Nettles R, Fowler Jr VG, Ryan T, et al. *Proposed modifications to the Duke criteria for the diagnosis of infective endocarditis*. Clinical Infectious Diseases 2000;30: 633–8.
57. Raad I. *Intravascular-catheter-related infections*. Lancet 1998;351: 893–8.
58. Messing B, Man F, Colimon R, Thuillier F, Beliah M, et al. *Antibiotic lock technique is an effective treatment of bacterial catheter related sepsis during parenteral nutrition*. Clinical Nutrition 1990;9: 220–7.
59. Randolph AG, Cook DJ, Gonzales CA, Andrew M. *Benefit of heparin in peripheral venous and arterial catheters: systematic review and meta-analysis of randomised controlled trials*. British Medical Journal 1998; 316: 969–75.
60. Goode CJ, Titler M, Rakel B, et al. *A meta-analysis of effects of heparin flush and saline flush: quality and cost implications*. Nursing Research 1991;40:324–30.
61. Peterson FY, Kirchoff KT. *Analysis of the research about heparinized versus nonheparinized intravascular lines*. Heart and Lung 1991;20: 631–40.
62. Jacobs BR, Schilling S, Doellman D, Hutchinson N, Rickey M, Nelson S, et al. *Central venous catheter occlusion: a prospective, controlled trial examining the impact of a positive-pressure valve device*. Journal of Parenteral and Enteral Nutrition 2004; 28: 113–8.
63. Jay R, Grove JR, Pevec WC. *Venous thrombosis related to peripherally inserted central catheters*. Journal of Vascular and Interventional Radiology 2000;11: 837–40.
64. Ong B, Gibbs H, Catchpole I, Hetherington R, Harper J, et al. *Peripherally inserted central catheters and upper extremity deep vein thrombosis*. Australasian Radiology 2006; 50: 451–4.