

## 10. BIBLIOGRAFIA

1. Registered Nurses Association of Ontario (RNAO). *Screening for delirium, dementia and depression in older adult*. Toronto, Canada: Registered Nurses Association of Ontario (RNAO ); 2003 Nov. 88 p.
2. American Psychiatric Association. *Practice guideline for the treatment of patients with delirium*. 1999. Su [http://www.psych.org/clin\\_res/pg\\_delirium.html](http://www.psych.org/clin_res/pg_delirium.html) (data di ultima consultazione 30/04/2008)
3. Royal College of Physicians. *Guidelines for the prevention, diagnosis and management of delirium in older people in hospital*. 2005. Su <http://www.sign.ac.uk/guidelines/fulltext/50/section6.html> (data di ultima consultazione 30/04/2008)
4. Rubin FH. *Preoperative and postoperative issue in the orthopaedic surgical patient*. Operative techniques in Orthopaedics 2002; 112(2): 60-63.
5. Evidence based nursing *Assistenza al paziente con delirium. Schede informative 18 per il miglioramento dell' assistenza infermieristica-riabilitativa-ostetrica*. Su <http://evidencebasednursing.it> ( Data di ultima consultazione 30/04/2008)

6. Eden BM, Foreman MD. *Problems associated with underrecognition of delirium in critical care : a case study*. Heart Lung 1996; 25:388-400.
7. Edelstein DM, Aharonoff GB, Karp A ,Capla EL, Zukerman JD, Koval KJ. *Effect of Postoperative delirium on outcome after hip fracture*. Clinical orthopaedics and related research 2004; 422 :195-200
8. Victorian Government Department of Human Services. *Clinical practice guidelines for the management of delirium in older people*.Victorian Government Department of human Service .Melburne Australia 2006. Su <http://www.health.vic.gov.au/acute-agedcare> (Data di ultima consultazione 30/04/2008)
9. Truman B, Ely EW. *Monitoring delirium in critically ill patients. Using the confusion assesment method for the intensive care unit* . Critical Care Nurse 2003; 23 (2): 25-35
- 10.Schuurmans MJ, Duursma SA, Shortridge LM, Clevers GJ, Pel-Little R. *Elderly patients with a hip fracture :the risk for delirium*. Applied Nursing Research 2003 ;16 (2): 75-84
- 11.Olofsson B, Lundstrom M, Borssén B, Nyberg L, Gustafson Y. *Delirium is associated with poor rebilitathion outcomes in elderly patients treated for femoral neck fractures*. Scand J Caring Sci 2005; 19 :119-127
- 12.Zakriya KJ , Cristmas C , Wenz JF , Franckowiak S , Anderson R , Sieber EF. *Preoperative factors associated with postoperative*

- change in confusion assesment method scorein hip fracture patients. Anesth Analg 2002 ; 94 : 1628-32*
13. Marcantonio ER, Flacker JM, Wrigth J, Resnick NM. *Reducing delirium after hip fracture: a randomized trial . American Geriatric Society 2001 ; 49: 516-522*
14. Adunsky A, Levy R, Hiem M, Mizhai E, Arad M. *The unfavorable nature of postoperative delirium in elderly hip fracture patients. Arch. Gerontol. Geriatr. 2003 ;36 : 67-74*
15. Brauer C, Morrison RS, Silbezweig SB, Siu AL. *The cause of delirium in patients with hip fracture. Arch Med. 2000; 160: 1856-1860*
16. Università di Pisa , Facoltà di Medicina e Chirurgia . Vanni A. *Il delirium post-operatorio un problema di medicina perioperatoria. Tesi di Laurea.*
17. Inouye SK. *Delirium in older person. N Engl J Med 2006; 354: 1157-1165*
18. Mussi C, Savioli G. *Linee guida per la diagnosi e la terapia nel delirium nell' anziano. Giorn Gerontol . 2000 ; 48: 434-440*
19. Rockwood K. *The occurence and duration of syntoms in elderly patients with delirium . J Gerontol 1993 ; 48: M162-M166*
20. Mussi C. *Come si diagnostica . G. Gerontol 2004 ; 52: 368-375*
21. Cavazzuti F, Creminini G, Di Giulio P, Zanetti E. *Assistenza geriatrica oggi. Casa Editrice Ambrosiana . Milano 2008 ;18 : 186-189*

22. Forrest J , Willis L, Holm K, Kwon MS, Anderson MA, Foreman MD.  
*Recognizing quiet delirium* . AJN 2007; 107 (4 ) : 35-39
23. Inouye SK, Foreman MD, Mion LC, Katz KH, Cooney LM .*Nurses' recognition of Delirium and its simtoms. Comparison of Nurse and research rating* . Arch .Intern. Med . 2001 ; 161 : 2467-2473
24. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. Fourth edition. Washington: DC, American psychiatric association 1994
25. Jahnigan DW. *Delirium in the elderly hospitalized patients*. Hosp Pract 1990; 15: 135-157
26. World Health Organization. *The ICD Classification of Mental and Behavioural Disorders*. Geneva : Woeld Health Organization 1992
27. Edwards N. *Differentiating the three D's : Delirium ,Dementia, and Depression*. Medsurg Nursing 2003; 12(6) :347-357
28. Inouye SK, Van Dick CH, Alessi CA, Balkin S, Siegal AP, Horvitz RI. *Clarifyng confusion : the confusion assesment method* . Ann Intern Med 1990 ;113: 941-948
29. Inouye SK. *The confusion assesment method (CAM): training manual and coding guide* . 2003 Yale University School of Medicine
30. Neelon VJ, Champagne MT, Charlson JR, Funk SG. *The neecham confusion scale : costruction, validation and clinical testing*. Nursing Research 1996; 45: 324-330
31. Van Germet LA, Schuurmans MJ . *The eecham confusion scale and the delirium observatiene screening scale : capacity to*

- discriminate and ease of use in clinical practice* . BMC Nursing  
2007 ; 6:3
32. Johansson IS, Harmin EKF, Larsson G. *Psychometry testing of the neecham confusion scale among patients with hip fracture*.  
Research in Nursing & Health 2002 ; 25 : 203-211
33. Dupplins GS, Wikblad K. *Delirium : behavioural changes bifore and durino the podromale phase*. Journal of Clinical Nursing  
2004;13:609-616
34. Inouye SK, Viscoli CM, Horwitz RI, Hurst Ld, Tinetti ME. *A predictive model for delirium in hospitalized elderly medical patients based on admission characteristis*. Ann Intern Med 1993 ; 119: 474-481
35. Bergamini L, Bergamasco B, Mutani R. *Manuale di neurologia clinica*. Edizioni Libreria Cortina Torino 2001. 89-93
36. Inouye SK, Borgadus ST, Williams CS, Leo-Summers L, Agostini JV. *The role of adherence on the effectiveness of nonpharmacologic intervention* . Arch Intern Med. 2003; 163: 958-964
37. U.S. Food and Drug Administration. *Information for healthcare professionals haloperidol (marked as Haldol, Haldol Decanoate and Haldol Lactate)*. Departement of Health and Human Services 2007 .  
Su <http://www.fda.gov/cder/whatsnew.htm> (Data di ultima consultazione 15/04/2008)

38. Milisen K, Foreman MD, Wounters B, Driesen R, Godderis J, Abraham IL, Broos PL. *Documentation of delirium in elderly patients with hip fracture*. Journal of Gerontological Nursing 2002; 28 (11): 23-29
39. Gillins AJ, Mac Donald B. *Unmasking delirium. Nurse can play a key role in the prevention, detection and treatment of delirium in older adults* . American Journal of Nursing 2006 ; 102 (9) :
40. Milisen K, Foreman MD , Abraham IL , De Geest S , Godderis J, Vandermeulem E. *A nurse- led interdisciplinary intervention program for Delirium in elderly hip fracture patients*. J Am Geriatr Soc 2001; 49: 523-522
41. Inouye SK, Zhan H, Jones RN, Kiely DK, Yang F, Marcantonio E. *Risck factors for delirium at discharge. Development and Validation of a predictive model*. Arc Intern Med 2007; 167(13): 1406-1413
42. McAvay G, Van Ness PH, Bogardus ST, Zhang Y, Leslie D, Inouye SK. *Older adults discharged from the hospital with delirium:1-year outcomes*. JAGS 2006; 54 : 1245-1250