

BIBLIOGRAFIA

¹ Gensini GF, Zaninelli A, Bignamini AA, Ricci S, Segà R, Gandolfo C et al. SPREAD V edizione. Milano: Pubblicazioni Catel – Hyperphar Group SpA, 2007.

² Guidetti D, Raganini G, Peragallo M, Mosetti F, Lavallo T. Dossier InFad – anno 3, n. 40. Milano, Zadig, maggio 2008.

³ Ibidem.

⁴ Basaglia N. Progettare la Riabilitazione. Milano, Ermes, marzo 2002.

⁵ Cotran RS, Kumar V, Tucker C. Le basi patologiche delle malattie VI edizione. Padova, Piccin Nuova Libreria S.p.A. 2000: 1506-1515.

⁶ Ibidem.

⁷ Ibidem.

⁸ Opera già citata: (1).

⁹ Feigin VL, Lawes CM, Bennett DA, Anderson CS. Stroke epidemiology: a review of population-based studies of incidence, prevalence, and case-fatality in the late 20th century. *Lancet Neurology* 2003; 2: 43-53.

The Italian Longitudinal Study on Aging Working Group. Prevalence of chronic diseases in older Italians: comparing self-reported and clinical diagnoses. *J Epidemiol* 1997; 26: 995-1002.

¹⁰ Ricci S, Celani MG, La Rosa F, Vitali R, Duca E, Ferraguzzi R et al. SEPIVAC: a community-based study of stroke incidence in Umbria, Italy. *J Neurol Neurosurg Psychiatry* 1991; 54: 695-698.

¹¹ D'Alessandro G, Di Giovanni M, Roveyaz L, Iannizzi L, Compagnoni MP, Blanc S et al. Incidence and prognosis of stroke in the Valle d'Aosta, Italy: first-year result of a community-based study. *Stroke* 1992; 23: 1712-1715.

¹² Lauria G, Gentile M, Fassetta G, Casetta I, Agnoli F, Andreotta G et al. Incidence and prognosis of stroke in the Belluno province, Italy. First-year results of a community-based study. *Stroke* 1995; 20:1787-1793.

¹³ D'Alessandro G, Bottacchi E, Di Giovanni M, Martinazzo C, Sironi L, Lia C, Carenini L et al. Temporal trends of stroke in Valle d'Aosta, Italy. Incidence and 30-day fatality rates. *Neurol Sci* 2000; 21:13-18.

¹⁴ Di Carlo A, Inzitari D, Galati F, Baldereschi M, Giunta V, Grillo G et al. A prospective community-based study of stroke in Southern Italy: the Vibo Valentia incidence of stroke study (VISS). Methodology, incidence and case fatality at 28 days, 3 and 12 months. *Cerebrovasc Dis* 2003; 35: 410-417.

¹⁵ Di Carlo A, Launer LJ, Breteler MMB, Fratiglioni L, Lobo A, Martinez-Lage J et al. for the ILSA Working Group and the Neurologic Diseases in the Elderly Research Group. Frequency of stroke in Europe: A collaborative study of population-based cohorts. *Neurology* 2000; 16 (sup. 5): 28-33.

¹⁶ Sudlow CL, Warlow CP. Comparable studies of the incidence of stroke and its pathological types: results from an international collaboration. International Stroke Incidence Collaboration. *Stroke* 1997; 27: 491-499.

¹⁷ Opera già citata: (9).

¹⁸ Opera già citata: (6).

¹⁹ Pitacco G. Il modello di analisi della complessità assistenziale. *L'Infermiere* 2003; 8-9 (sup. 7): 22-27.

²⁰ *Ibidem*: 24 - grafico 1.

²¹ *Ibidem*: 26 - grafico 5.

²² *Ibidem*: 27 - grafico 6.

²³ Opera già citata: (1) 403 - figura 14-1.

²⁴ Cisari C, Diana G, Freddi A, Neri G, Pace P, Piscaglia MG. Linee organizzative per il governo clinico dello stroke. Documento interdisciplinare 2001.

²⁵ www.digilander.libero.it/aiol/nursing/accertamento.htm (u.c. 16/09/08).

²⁶ Silvestro A. Premessa. *L'Infermiere* 2003; 2/032' (sup. 6): 3.

Santullo A, Chiari P. Il valore delle scale. *L'Infermiere* 2003; 2/032' (sup. 6): 4-12.

Mosci D, Chiari P. Scala di Conley: uno strumento per la predizione del rischio di caduta del paziente anziano. *L'Infermiere* 2003; 2/032' (sup. 6): 17-20.

Bellingeri A. La valutazione del rischio di contrarre lesioni da pressione. *L'Infermiere* 2003; 2/032' (sup. 6): 21-25.

Santullo A. Scala Adl di Katz: uno strumento per la valutazione soggettiva dello stato funzionale. *L'Infermiere* 2003; 2/032' (sup. 6): 26-28.

Lumini E. Le scale di valutazione della funzione neurologica. *L'Infermiere* 2003; 2/032' (sup. 6): 29-32.

Opera già citata: (1)

²⁷ Rodgers H, Atkinson C, Bond. Randomized Controlled Trial of a Comprehensive Stroke Education Program for Patients and Caregivers. *Stroke* 1999; 30: 2585-2591.

Mant J, Carter J, Wade DT, Winner S. The impact of an information pack on patient with stroke and their carers: a randomized controlled trial. *Clinical rehabilitation*, 1998; 19: 465-76.

Chemerinski E, Robinson R, Arndt S, Kosier J. The effect of remission of post-stroke depression on activities of daily living in a double blinded randomized treatment study. *Journal of Nervous and Mental Disease*, 2001. 191: 421-425.

Opera già citata: (1).

Clark MS, Rubenach S, Winsor A. A randomized controlled trial of an education and counselling intervention for families after stroke. *Clin Rehabil*. November 2003. 19: 1242-1249.

Sulch D, Perez I, Melbourn A, Lalit K. Randomized controlled of integrated (managed) care pathway for stroke rehabilitation. *Clin Rehabil*. May 2000. 31: 1929-1934.

²⁸ Scottish Intercollegiate Guidelines Network (SIGN). Management of Patients with Stroke. Rehabilitation, Prevention and Management of complications, and Discharge Planning A. National clinical guideline, November 2002.

Page T, Lockwood C, Evans D. The prevention and management of shoulder pain in the hemiplegic patient. Blackwell Publishing. 2003. 7: 149-166.

www.continence-fdn.ca/indexeng.htm (u.c. 16/09/08).

Opera già citata: (1).

²⁹ Barere D. The natural history and functional consequences of dysphagia after hemispheric stroke. *Journal Neurosurg Psychiatry*; 1989. 4: 236-241.

Barri MA. Objective screening tools to assess cognitive impairment and depression. *Topics in Geriatric Rehabilitation*. 2002. 18: 28-46.

www.aafp.org/afp/980101ap/broderick.html (u.c. 20/09/2008).

Opera già citata: (1).

Basso A. *Conoscere e rieducare l'afasia*. Milano: Pensiero Scientifico, 2005.

Brugnolli A, Saiani L. *Piani di assistenza in medicina e chirurgia II edizione italiana*. Perugia: Idelson-Gnocchi, 2008.

Carpenito ML. *Diagnosi infermieristiche. Applicazione alla pratica clinica*. Roma: CEA, 2006.